

Embedding of the Kerr-Newman Black Hole Surface in Euclidean Space

Valeri P. Frolov*

¹*Theoretical Physics Institute, University of Alberta, Edmonton, AB, Canada, T6G 2J1*

(Dated: February 7, 2008)

We obtain a global embedding of the surface of a rapidly rotating Kerr-Newman black hole in an Euclidean 4-dimensional space.

PACS numbers: 04.70.Bw, 04.50.+h, 04.20.Jb

Alberta-Thy-01-06

I. INTRODUCTION

In this paper we discuss the problem of isometric embedding of the surface of a rapidly rotating black hole in a flat space.

It is well known that intrinsically defined Riemannian manifolds can be isometrically embedded in a flat space. According to the Cartan-Janet [1, 2] theorem, every analytic Riemannian manifold of dimension n can be locally real analytically isometrically embedded into $\mathbb{E}^N$ with $N = n(n+1)/2$. The so called Fundamental Theorem of Riemannian geometry (Nash, 1956 [3]) states that every smooth Riemannian manifold of dimension n can be globally isometrically embedded in a Euclidean space $\mathbb{E}^N$ with $N = (n+2)(n+3)/2$.

The problem of isometric embedding of 2D manifolds in $\mathbb{E}^3$ is well studied. It is known that any compact surface embedded isometrically in $\mathbb{E}^3$ has at least one point of positive Gauss curvature. Any 2D compact surface with positive Gauss curvature is always isometrically embeddable in $\mathbb{E}^3$, and this embedding is unique up to rigid rotations. (For general discussion of these results and for further references, see e.g. [4]). It is possible to construct examples when a smooth geometry on a 2D ball with negative Gauss curvature cannot be isometrically embedded in $\mathbb{E}^3$ (see e.g. [5, 6]). On the other hand, it is easy to construct an example of a global smooth isometric embedding for a surface of the topology S^2 which has both, positive and negative Gauss curvature ball-regions, separated by a closed loop where the Gauss curvature vanishes. An example of such an embedding is shown in Figure 1 [10].

The surface geometry of a charged rotating black hole and its isometric embedding in $\mathbb{E}^3$ was studied long time ago by Smarr [7]. He showed that when the dimensionless rotation parameter $\alpha = J/M^2$ is sufficiently large, there are two regions near poles of the horizon surface where the Gauss curvature becomes negative. Smarr proved that these regions cannot be isometrically embedded (even locally) in $\mathbb{E}^3$ as a revolution surface, but such local embedding is possible in a 3D Minkowsky space. More recently different aspects of the embedding of a surface of a rotating black hole and its ergosphere in $\mathbb{E}^3$




FIG. 1: This picture shows the "croissant" surface. A solid line separates two regions with opposite signs of the Gauss curvature. Each of these regions has the topology of a 2D ball. The Gauss curvature is negative in the upper ball-region.

were discussed in [8, 9]. A numerical scheme for construction of the isometric embedding for surfaces with spherical topology was proposed in [9]. The surface geometry of a rotating black hole in an external magnetic field and its embedding in $\mathbb{E}^3$ was studied in [12, 13, 14].

The purpose of this paper is to obtain the global isometric embedding of a surface of a rapidly rotating black hole in $\mathbb{E}^4$. In Section 2 we discuss general properties of 2D axisymmetric metrics and prove that if the Gauss curvature is negative at the fixed points of the rotation group it is impossible to isometrically embed a region containing such a fixed point in $\mathbb{E}^3$. In Section 3 we demonstrate that such surfaces can be globally embedded in $\mathbb{E}^4$. We obtain the embedding of surfaces of rapidly rotating black holes in $\mathbb{E}^4$ in an explicit form in Section 4. Section 5 contains a brief summary and discussions.

II. GEOMETRY OF 2D AXISYMMETRIC DISTORTED SPHERES

Let us consider an axisymmetric deformation S of a unit sphere S^2 . Its metric can be written in the form

$$dl^2 = h(x)dx^2 + f(x)d\phi^2. \quad (1)$$

Here $\xi = \partial_\phi$ is a Killing vector field with closed trajectories. Introducing a new coordinate $\mu = \int dx \sqrt{h f}$ one can rewrite (1) in the form

$$dl^2 = f(\mu)^{-1}d\mu^2 + f(\mu)d\phi^2. \quad (2)$$

We assume that the function f is positive inside the interval (μ_0, μ_1) and vanishes at its ends. We choose $\mu_0 = -1$. The surface area of S is $2\pi(\mu_1 + 1)$. By multiplying the

*Electronic address: frolov@phys.ualberta.ca

metric (1) by a constant scale factor $(\mu_1 + 1)^{-1/2}$ one can always put $\mu_1 = 1$. We shall use this choice, for which the surface area of S is 4π and the fixed points of ξ are located at $\mu = \pm 1$.

The metric (2) is regular (no conical singularities) at the points $\mu = \pm 1$ (where $r = 0$) if $f'(\pm 1) = \mp 2$. (Here and later $(\dots)' = d(\dots)/dz$.) The Gaussian curvature for the metric (2) is

$$K = -\frac{1}{2}f'' . \quad (3)$$

Let us introduce a new coordinate $r = \sqrt{f(\mu)}$. The metric (2) in these coordinates is

$$dl^2 = V(r)dr^2 + r^2 d\phi^2 , \quad V = \frac{4}{f'^2} . \quad (4)$$

We denote by K_0 the value of K at a fixed point of rotations. Then in the vicinity of this point one has

$$V \approx 1 + \frac{1}{2}K_0 r^2 + \dots . \quad (5)$$

For $K_0 < 0$ the region in the vicinity of the fixed point cannot be embedded as a revolution surface in a Euclidean space $\mathbb{E}^n$ for any $n \geq 3$. Indeed, consider a space $\mathbb{E}^n$ with the metric

$$dS^2 = dX^2 + dY^2 + \sum_{i=3}^n dZ_i^2 . \quad (6)$$

For the surface of revolution $X = r \cos \phi$, $Y = r \sin \phi$, $Z_i = Z_i(r)$ the induced metric is

$$dl^2 = V(r)dr^2 + r^2 d\phi^2 , \quad V(r) = 1 + \sum_{i=3}^n (dZ_i/dr)^2 . \quad (7)$$

For a regular surface $V(0) = 1$ and $V(r) \geq 1$ in the vicinity of $r = 0$. According to (5) this is impossible when $K_0 < 0$.

We show now that if $K_0 < 0$ then a ball-region near the fixed point p_0 of axisymmetric 2D geometry cannot be isometrically embedded in $\mathbb{E}^3$. Let us assume that such an embedding (not necessarily as a revolution surface) exists. One can choose coordinates (X^1, X^2, Z) in $\mathbb{E}^3$ so that $X^1 = X^2 = 0$ at p_0 , and in its vicinity

$$Z = \frac{1}{2}(k_1 X^{1^2} + k_2 X^{2^2}) + \dots , \quad (8)$$

where k_a ($a = 1, 2$) are principal curvatures at p_0 . Here and later ‘dots’ denote omitted higher order terms. The metric on this surface induced by its embedding is

$$dl^2 = (1 + k_1^2 X^{1^2})dX^{1^2} + (1 + k_2^2 X^{2^2})dX^{2^2} + 2k_1 k_2 X^1 X^2 dX^1 dX^2 + \dots . \quad (9)$$

In the vicinity of p_0 the Killing vector ξ generating rotations has the form

$$\xi = p^a \partial_a , \quad (10)$$

where p^a ($a = 1, 2$) are regular functions of (X^1, X^2) vanishing at $(0, 0)$. Their expansion near p_0 has the form

$$p^a = P_b^a X^b + P_{bc}^a X^b X^c + P_{bcd}^a X^b X^c X^d + \dots . \quad (11)$$

Consider the Taylor expansion near p_0 of the Killing equation

$$\xi_{a;b} = \xi_{a,b} - \Gamma_{ab}^c \xi_c = 0 \quad (12)$$

in the metric (9). Since the expansion of both Γ_{ab}^c and ξ_c starts with a linear in X^a terms, the equation (12) can be used to obtain restrictions on the coefficients P_b^a , P_{bc}^a , and P_{bcd}^a in (11). Simple calculations give

$$P_1^1 = P_2^2 = 0 , \quad P_2^1 = -P_1^2 = q , \quad P_{bc}^a = P_{bcd}^a = 0 , \quad (13)$$

$$qk_1(k_1 - k_2) = qk_2(k_1 - k_2) = 0 . \quad (14)$$

If the Killing vector does not vanish identically then $q \neq 0$ and the equations (14) imply that $k_1 = k_2$. This contradicts to the assumption of the existence of the embedding with $K_0 = k_1 k_2 < 0$.

III. EMBEDDING OF A 2D SURFACE WITH $K_0 < 0$ IN $\mathbb{E}^4$

Increasing the number of dimensions of the flat space from 3 to 4 makes it possible to find an isometric embedding of 2D manifolds with $K_0 < 0$. Denote by (X, Y, Z, R) Cartesian coordinates in $\mathbb{E}^4$ and determine the embedding by equations

$$X = \frac{r}{\Phi_0} \xi(\psi) , \quad Y = \frac{r}{\Phi_0} \eta(\psi) , \quad Z = \frac{r}{\Phi_0} \zeta(\psi) , \quad (15)$$

$$R = R(r) , \quad (16)$$

where $0 \leq \psi \leq 2\pi$, and functions ξ , η and ζ obey the condition

$$\xi^2(\psi) + \eta^2(\psi) + \zeta^2(\psi) = 1 . \quad (17)$$

In other words, $\mathbf{n} = (\xi, \eta, \zeta)$ as a function of ψ is a line on a unit sphere S^2 . We require that this line is a smooth closed loop ($\mathbf{n}(0) = \mathbf{n}(2\pi)$) without self-intersections. Since a loop on a unit sphere allows continuous deformations preserving its length, there is an ambiguity in the choice of functions (ξ, η, ζ) .

We denote $\Phi = (\xi_{,\psi}^2 + \eta_{,\psi}^2 + \zeta_{,\psi}^2)^{1/2}$ then

$$2\pi\Phi_0 = \int_0^{2\pi} d\psi \Phi(\psi) . \quad (18)$$



FIG. 2: The surface shown at this picture is formed by straight lines passing through a point $r = 0$. Its Gauss curvature vanishes. The surface has a cone singularity at $r = 0$ with a negative angle deficit.

is the length of the loop. Instead of the coordinate ψ it is convenient to use a new angle coordinate ϕ which is proportional to the proper length of a curve $r = \text{const}$

$$\phi = \Phi_0^{-1} \int_0^\psi d\psi' \Phi(\psi'). \quad (19)$$

The coordinate ϕ is a monotonic function of ψ and for $\psi = 0$ and $\psi = 2\pi$ it takes values 0 and 2π , respectively.

Equations (15) give the embedding in $\mathbb{E}^3$ of a linear surface formed by straight lines passing through $r = 0$. This surface has $K = 0$ outside the point $r = 0$ where, in a general case, it has a cone-like singularity with the angle deficit $2\pi(1 - \Phi_0)$.

We shall use the embedding (15)–(16) for the case when the angle deficit is negative. In this case one can use, for example, the following set of functions

$$\xi = \cos \psi / F, \quad \eta = \sin \psi / F, \quad \zeta = a \sin(2\psi) / F, \quad (20)$$

$$F = \sqrt{1 + a^2 x^2}, \quad x = \sin^2(2\psi). \quad (21)$$

This embedding for the functions (ξ, η, ζ) defined by (20)–(21) is shown in Figure 2.

For this choice

$$\Phi = (1 + 4a^2 - 3a^2 x)^{1/2} (a^2 x + 1)^{-1}, \quad (22)$$

$$\Phi_0 = \frac{1}{\pi} \int_0^1 \frac{dx(1 + 4a^2 - 3a^2 x)^{1/2}}{\sqrt{x(1-x)(a^2 x + 1)}}. \quad (23)$$

Calculations give

$$\Phi_0 = \frac{8}{\pi \sqrt{1 + 4a^2}} [(1 + a^2)\Pi(-a^2, k) - 3/4K(k)], \quad (24)$$

$$k = \sqrt{3}a / (1 + 4a^2). \quad (25)$$

Here $K(k)$ and $\Pi(\nu, k)$ are complete elliptic integrals of the first and third kind, respectively. The function Φ_0




FIG. 3: Φ_0 as a function of the parameter a .

monotonically increases from 1 (at $a = 0$) to 2 (at $a \rightarrow \infty$) (see Fig. 3).

The induced metric for the embedded 2D surface defined by (15)–(16) is

$$dl^2 = [\Phi_0^{-2} + (dR/dr)^2] dr^2 + r^2 d\phi^2. \quad (26)$$

If the angle deficit is positive ($\Phi_0 < 1$), the pole point $r = 0$ in the metric (15) remains a cone singular point for any $R(r)$. For $\Phi > 1$ (the negative angle deficit), the pole-point $r = 0$ in the metric (26) is regular if $(dR/dr)_0^2 = 1 - \Phi_0^2$. By comparing (4) and (26) one obtains

$$r = f^{1/2}, \quad (dR/dr)^2 = (V - \Phi_0^{-2}). \quad (27)$$

This relation gives the following equation relating $R(\mu)$ with $f(\mu)$

$$R' = (1 - f'^2 / (4\Phi_0^2))^{1/2} f^{-1/2}. \quad (28)$$

It is easy to check that $R'' = 0$ at points where $f' = 0$. In order R' to be real, the following condition must be valid $\Phi_0 \geq \frac{1}{2} \max_{\mu \in (-1, 1)} |f'(\mu)|$. At a point where $|f'|$ reaches its maximum the quantity $f'' = -2K$ vanishes. Thus it is sufficient to require that Φ_0 is greater or equal to the values of $|f'|$ calculated at the points separating regions with the positive and negative Gauss curvature.

IV. EMBEDDING OF THE SURFACE OF THE KERR-NEWMAN HORIZON IN $\mathbb{E}^4$

The surface geometry of the Kerr-Newman black hole is described by the metric $ds^2 = N^2 dl^2$, where

$$dl^2 = (1 - \beta^2 \sin^2 \theta) d\theta^2 + \sin^2 \theta [1 - \beta^2 \sin^2 \theta]^{-1} d\phi^2, \quad (29)$$

$$N = (r_+^2 + a^2)^{1/2}, \quad \beta = a(r_+^2 + a^2)^{-1/2}. \quad (30)$$

Here $0 \leq \theta \leq \pi$ and $0 \leq \phi \leq 2\pi$. The metric dl^2 is normalized so that the area of the surface with this metric is 4π . In the coordinates $\mu = \cos \theta$ the metric (29) takes the form (2) with

$$f(\mu) = (1 - \mu^2)[1 - \beta^2(1 - \mu^2)]^{-1}. \quad (31)$$




FIG. 4: Plot for R as a function of μ for $\beta = 0.7$.

For the black hole with mass M , charge Q and the angular momentum $J = Ma$

$$r_+ = M - (M^2 - a^2 - Q^2)^{1/2}. \quad (32)$$

The rotation parameter a and mass M can be written in terms of the distortion parameter β as follows

$$a = \beta N, \quad M = \frac{1}{2}N(1 - \beta^2)^{-1/2}(1 + Q^2/N^2). \quad (33)$$

The condition $M^2 \geq a^2 + Q^2$ for given parameters N and β requires that [7]

$$0 \leq Q \leq N(1 - \beta^2)^{1/2}, \quad (34)$$

$$\frac{1}{2}N(1 - \beta^2)^{-1/2} \leq M \leq N(1 - \beta^2)^{1/2}. \quad (35)$$

The distortion parameter has its maximal value $\beta_{max} = 1/\sqrt{2}$ for $Q = 0$. The Gauss curvature of the surface with the metric (29) is

$$K = [1 - \beta^2(1 + 3\mu^2)][1 - \beta^2(1 - \mu^2)]^{-3}. \quad (36)$$

For $\frac{1}{2} < \beta \leq \frac{1}{\sqrt{2}}$ the Gauss curvature is negative in the vicinity of poles in the region $\mu_c \leq |\mu| \leq 1$

$$\mu_c = (1 - \beta^2)^{1/2}(\sqrt{3}\beta)^{-1}. \quad (37)$$

At $|\mu| = \mu_c$ the Gauss curvature vanishes. As it was shown earlier, at this point $|f'|$ has its maximum

$$|f'|_{\max} = |f'|_{\mu_c} = \frac{3\sqrt{3}}{8\beta(1 - \beta^2)^{3/2}}, \quad (38)$$

and one must choose the parameter Φ_0 so that $\Phi_0 \geq \frac{1}{2}|f'|_{\max}$. Simplest possible choice is

$$\Phi_0 = \frac{1}{2}|f'|_{\mu_c}. \quad (39)$$

Using (39) and integrating the equation (28) one determines R as a function of μ . A plot of this function for $\beta = 0.7$ is shown in Figure 4. Plot 1 at Figure 5 shows R as a function of r for the same values of β .




FIG. 5: Plot for R as a function of r (curve 1) and ρ (curve 2) for $\beta = 0.7$.




FIG. 6: By gluing these two figures along their edges one obtains a $2D$ surface without angle deficits and isometric to the surface of a rotating black hole ($\beta = 0.7$.)

The metric (26) can also be written in the form

$$dl^2 = [1 + (dR/d\rho)^2]d\rho^2 + \Phi_0^2\rho^2d\phi^2, \quad (40)$$

where $\rho = r/\Phi_0$. Plot 2 at Figure 5 shows R as a function of ρ for $\beta = 0.7$. The metric (40) coincides locally with the metric on the revolution surface determined by the equation $R = R(\rho)$ in $\mathbb{E}^3$. This does not give a global isometric embedding since the period of the angle coordinate is $2\pi\Phi_0$. This surface can be obtained by gluing two figures shown in Figure 6 along their edges. For the left figure ϕ changes from 0 to π , while for the right one it changes from π to $2\pi\Phi_0$.

V. CONCLUDING REMARKS

We demonstrated that a surface of a rapidly rotating black hole, which cannot be isometrically embedded in E^3 , allows such a global embedding in E^4 . To construct this embedding one considers first a $2D$ surface in E^3 formed by straight lines passing through one point ($r=0$) which has a cone singularity at $r = 0$ with negative angle deficit. Its Gauss curvature outside $r = 0$ vanishes. Next element of the construction is finding a function $R(r)$. The revolution surface for this function in E^3 has a positive angle deficit at $r = 0$. By combining these two maps in such a way that positive and negative angle deficits cancel one another, one obtains a regular global embedding in E^4 . This construction can easily be used

to find the embedding in E^4 of surfaces of rapidly rotating stationary black holes distorted by an action of external forces or fields, provided the axial symmetry of the spacetime is preserved. An interesting example is a case of a rotating black hole in a homogeneous at infinity magnetic field directed along the axis of the rotation (see e.g. [12, 13, 14].

Acknowledgments

This work was supported by the Natural Sciences and Engineering Research Council of Canada and by the Kil-lam Trust.

-
- [1] M. Janet, Ann. Soc. Polon. Math. **5**, 38 (1926).
 - [2] E. Cartan, Ann. Soc. Polon. Math. **6**, 1 (1927).
 - [3] J. Nash, Annals of Math. **63**, 20 (1956).
 - [4] M. Berger, *A Panoramic View of Riemannian Geometry*, Springer-Verlag, Berlin-Heidelberg-New York, (2003).
 - [5] A. V. Pogorelov, Dokl. Akad. Nauk SSSR, **198**, 42 (1971); English translation in Soviet Math. Dokl. **12**, 729 (1971).
 - [6] N. Nadirashvili and Y. Yuan, *Counterexamples for Local Isometric Embedding*, Eprint math.dg/0208127.
 - [7] L. Smarr, Phys. Rev. **D7**, 289 (1973).
 - [8] N. Pelavas, N. Neary, and K. Lake, Class. Quant. Grav. **18**, 1319 (2001).
 - [9] M. Bondarescu, M. Alcubierre, and E. Seidel, Class. Quant. Grav. **19**, 375 (2002).
 - [10] This example demonstrates that the Proposition 4.1 of the paper [11] is wrong.
 - [11] J. Stryla, Class. Quant. Grav. **15**, 2303 (1998).
 - [12] W. J. Wild and R. M. Kerns, Phys. Rev. **D21**, 332 (1980).
 - [13] W. J. Wild, R. M. Kerns, and W. F. Drish, Jr., Phys. Rev. **D23**, 829 (1981).
 - [14] R. Kulkarni and N. Dadhich, Phys. Rev. **D33**, 2780 (1986).